

More on Tanaka-Tagoshi Parametrization of post-1PN Spin-Free Gravitational Wave Chirps: Equispaced and Cardinal Interpolated Lattices

R.P. Croce and Th. Demma
Wavesgroup, D.I.³E., University of Salerno, Italy

V. Pierro and I.M. Pinto
Wavesgroup, University of Sannio at Benevento, Italy
(Dated: February 7, 2008)

The conclusions obtained in gr-qc/0101067 are shown to be valid also if the full 2.5PN expansion of the chirp phase is used.

PACS numbers: 04.80.Nn, 95.55.Ym, 95.75.Pq, 97.80.Af

In a recent paper [1] we shew that the spin-free gravitational wave templates parametrization introduced by Tanaka and Tagoshi in [2] is effective to set up *uniformly spaced* post-1PN template lattices subject to a given minimal match constraint. Lattice-uniformity makes cardinal interpolation of the match [3], [4] in the 2D Tagoshi-Tanaka parameter space straightforward, yielding a reduction in the 2PN template density and total number by a factor ~ 4 at $\Gamma = 0.97$.

In [1] the 2PN approximant of the (spectral, spin-free) chirp phase has been used. On the other hand, in [2] the 2.5PN phase (originated in [5]) has been used, which contains (besides an irrelevant f -independent term, which is absorbed in the unknown coalescency phase ϕ_C) the additional term:

$$\frac{\pi}{128\eta} \left(\frac{38645}{252} + 5\eta \right) \cdot \log(f/f_0), \quad (0.1)$$

where $\eta = m_1 m_2 / m^2$ is the symmetric mass-ratio, and f_0 is a suitable scaling frequency [6]. It has been pointed out [5] that the 2.5PN phase agrees, in the limit $\eta \rightarrow 0$, with the one obtained by Tagoshi and Sasaki in the (perturbative) test mass-limit solution of the spin-free relativistic two-body problem [7]. While this might be a good reason to include the 2.5PN term (0.1) in the template phase, it does *not* result into better overlaps with exact (numerically computed) waveforms. Indeed, as shown in [8], 2.5PN templates yield generally *poorer* overlaps (and larger biases) as compared to 2PN, in view of the peculiar (oscillating) behaviour of standard PN approximants [8].

However, the conceptual question is posed whether the results in [1] still hold if the 2.5PN logarithmic term (0.1) is included in the template phase. The purpose of this note is to show that the main conclusions in [1] do apply in this case too, though a number of differences are noted, which are summarized below.

The major difference, after the inclusion of the 2.5PN logarithmic term (0.1) in the phase, is a moderate increase in the area of the Tanaka-Tagoshi (flattened) manifold \mathcal{T} , yielding a correspondingly larger number of templates, as seen by comparing Fig. 1 and Table I of this note to Fig. 3 and Table III of [1], respectively.

The gaussian curvature of the spin-free manifold retains the same order of magnitude, as seen by comparing Fig. 2 of this note to Figs 1 and 2 of [1].

The main relevant conclusions in [1] do *not* change. As shown in [1], the minimal match degradation due to using the (flattened) Tanaka-Tagoshi (spin-free) parameter-space manifold in place of the (curved) true one, can be gauged in terms of the quantity η defined in Eq. (III.6) of [1]. Even using the 2.5PN phase, the quantity η is still of the same order of magnitude as the corresponding one computed in [1]. This is seen by comparing Fig. 3 of this note to Figs 5 and 6 of [1].

Acknowledgements

The Authors thank dr. A. Pai (IUCAA) for having posed the problem, and dr. B. Owen (Univ. of Wisconsin Milwaukee) for pinpointing an error in the first version of this note.

-
- [1] R.P. Croce, Th. Demma, V. Pierro and I.M. Pinto, gr-qc/0101067; in print on Phys. Rev. **D63** (2001).
 - [2] T. Tanaka and H. Tagoshi, Phys. Rev. **D62**, 082001 (2000).
 - [3] R.P. Croce, Th. Demma, V. Pierro and I.M. Pinto, Phys. Rev. **D62**, 124020 (2000).

- [4] R.P. Croce, Th. Demma, V. Pierro, I.M. Pinto, D. Churches and B.S. Sathyaprakash, Phys. Rev. **D62**, 121101(R) (2000).
- [5] L. Blanchet, T. Damour, B. Iyer, C.M. Will and A. Wiseman, Phys. Rev. Lett., **74**, 3515 (1995).
- [6] Scaling f to f_0 in the argument of (0.1) entails an irrelevant constant additive phase which is absorbed in the unknown coalescence phase ϕ_C .
- [7] H. Tagoshi and M. Sasaki, Progr. Th. Phys., **92**, 745 (1994).
- [8] T. Damour, B. Iyer and B.S. Sathyaprakash, Phys. Rev. **D57**, 885 (1998).

Antenna	Simplex area [sec^2]	No. of templates at $\Gamma = 0.97$
TAMA300	10266	$1.71 \cdot 10^5$
GEO600	57093	$9.51 \cdot 10^5$
LIGO-I	24732	$4.12 \cdot 10^5$
VIRGO	776392	$1.29 \cdot 10^7$

Table I - 2.5PN flat simplex area and number of templates at $\Gamma = .97$,
for $0.2M_\odot \leq m_1 \leq m_2 \leq 10M_\odot$.




Fig 1 – Left: the 2.5PN flat simplexes corresponding to $0.2 M_{\odot} \leq m_1 \leq m_2 \leq 10 M_{\odot}$ for TAMA300, GEO600, LIGO-I and VIRGO; right: the euclidean distance between the 2.5PN manifolds \mathcal{P} and \mathcal{T} as a function of (x_1, x_2) for TAMA300, GEO600, LIGO-I and VIRGO.





Fig. 2 – Left: gaussian curvature K [sec^{-2}] of spin-free 2.5PN parameter space manifold vs. m_1, m_2 for LIGO-1; right: the same quantity vs. $m_1=m_2$ (worst case) for TAMA300, GEO600, LIGO-1 and VIRGO.





Fig 3 – Left: the 2.5PN quantity η (eq.III.6 of [1]) vs. m_1, m_2 for LIGO-I; right: the same quantity vs. $m_1=m_2$ (worst case) for TAMA300, GEO600, LIGO-I and VIRGO.